

# **ORF522 – Linear and Nonlinear Optimization**

## **2. Linear optimization**

**Bartolomeo Stellato – Fall 2022**

# Today's agenda

Readings: [Chapter 1, Bertsimas, Tsitsiklis]

- Linear optimization in inner-product and matrix notation
- Optimization terminology
- Standard form
- Piecewise-linear minimization
- Examples

# Where does linear optimization appear?

Supply chain management

Assignment problems

Scheduling and routing problems

Finance

Optimal control problems

Network design and network operations

Many other domains...

# Vector notations

By default, all vectors are column vectors and denoted by

$$x = (x_1, \dots, x_n)$$

The transpose of a vector is  $x^T$

$a^T x$  is the inner product between  $a$  and  $x$

$$a^T x = a_1 x_1 + \cdots + a_n x_n = \sum_{i=1}^n a_i x_i$$

# Linear optimization

## Linear Programming (LP)

minimize  $\sum_{i=1}^n c_i x_i$

subject to  $\sum_{j=1}^n a_{ij} x_j \leq b_i, \quad i = 1, \dots, m$   
 $\sum_{j=1}^n d_{ij} x_j = f_i, \quad i = 1, \dots, p$

Objective function and constraints are **linear in the decision variables**

Belongs to **continuous optimization**

# Linear optimization

## Inner product notation

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^n c_i x_i \\ \text{subject to} & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad i = 1, \dots, m \\ & \sum_{j=1}^n d_{ij} x_j = f_i, \quad i = 1, \dots, p \end{array} \longrightarrow \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & a_i^T x \leq b_i, \quad i = 1, \dots, m \\ & d_i^T x = f_i, \quad i = 1, \dots, p \end{array}$$

$c, a_i, d_i$  are  $n$ -vectors

$$c = (c_1, \dots, c_n)$$

$$a_i = (a_{i1}, \dots, a_{in})$$

$$d_i = (d_{i1}, \dots, d_{in})$$

# Linear optimization

## Matrix notation

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^n c_i x_i \\ \text{subject to} & \sum_{j=1}^n a_{ij} x_j \leq b_i, \quad i = 1, \dots, m \\ & \sum_{j=1}^n d_{ij} x_j = f_i, \quad i = 1, \dots, p \end{array} \longrightarrow \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & Dx = f \end{array}$$

$A$  is  $m \times n$ -matrix with elements  $a_{ij}$  and rows  $a_i^T$

$D$  is  $p \times n$ -matrix with elements  $d_{ij}$  and rows  $d_i^T$

All (in)equalities are elementwise

# Optimization terminology

$$\begin{aligned} & \text{minimize} && c^T x \\ & \text{subject to} && Ax \leq b \\ & && Dx = f \end{aligned}$$

$x$  is **feasible** if it satisfies the constraints  $Ax \leq b$  and  $Dx = f$

The **feasible set** is the set of all feasible points

$x^*$  is **optimal** if it is feasible and  $c^T x^* \leq c^T x$  for all feasible  $x$

The **optimal value** is  $p^* = c^T x^*$

**Unbounded problem:**  $c^T x$  is unbounded below on the feasible set ( $p^* = -\infty$ )

**Infeasible problem:** feasible set is empty ( $p^* = +\infty$ )

# Standard form

## Definition

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

- Minimization
- Equality constraints
- Nonnegative variables

- Matrix notation for **theory**
- Standard form for **algorithms**

# Standard form

## Transformation tricks

### Change objective

If “maximize”, use  $-c$  instead of  $c$  and change to “minimize”.

### Eliminate inequality constraints

If  $Ax \leq b$ , define  $s$  and write  $Ax + s = b$ ,  $s \geq 0$ .

$s$  are the **slack variables**

If  $Ax \geq b$ , define  $s$  and write  $Ax - s = b$ ,  $s \geq 0$ .

### Change variable signs

If  $x_i \leq 0$ , define  $y_i = -x_i$ .

### Eliminate “free” variables

If  $x_i$  unconstrained, define  $x_i = x_i^+ - x_i^-$ , with  $x_i^+ \geq 0$  and  $x_i^- \geq 0$ .

# Standard form

## Transformation example

$$\begin{array}{ll} \text{minimize} & 2x_1 + 4x_2 \\ \text{subject to} & x_1 + x_2 \geq 3 \\ & 3x_1 + 2x_2 = 14 \\ & x_1 \geq 0 \end{array}$$



$$\begin{array}{ll} \text{minimize} & 2x_1 + 4x_2^+ - 4x_2^- \\ \text{subject to} & x_1 + x_2^+ - x_2^- - x_3 = 3 \\ & 3x_1 + 2x_2^+ - 2x_2^- = 14 \\ & x_1, x_2^+, x_2^-, x_3 \geq 0. \end{array}$$

# Linear, affine and convex functions

**Linear function:**  $f(x) = a^T x$

$$f(\alpha x + \beta y) = \alpha f(x) + \beta f(y), \quad \forall x, y \in \mathbf{R}^n, \alpha, \beta \in \mathbf{R}$$

**Affine function:**  $f(x) = a^T x + b$


$$f(\alpha x + (1 - \alpha)y) = \alpha f(x) + (1 - \alpha)f(y), \quad \forall x, y \in \mathbf{R}^n, \alpha \in \mathbf{R}$$

**Convex function:**

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y), \quad \forall x, y \in \mathbf{R}^n, \alpha \in [0, 1]$$

# Convex piecewise-linear functions

$$f(x) = \max_{i=1,\dots,m} (a_i^T x + b_i)$$



# Convex piecewise-linear minimization

**Equivalent linear optimization**

$$\text{minimize } f(x) = \max_{i=1,\dots,m} (a_i^T x + b_i)$$

$$\begin{aligned} & \text{minimize} && t \\ & \text{subject to} && a_i^T x + b_i \leq t, \quad i = 1, \dots, m \end{aligned}$$

**Matrix notation**

$$\begin{aligned} & \text{minimize} && \tilde{c}^T \tilde{x} \\ & \text{subject to} && \tilde{A} \tilde{x} \leq \tilde{b} \end{aligned}$$

$$\tilde{x} = \begin{bmatrix} x \\ t \end{bmatrix}, \quad \tilde{c} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad \tilde{A} = \begin{bmatrix} a_1^T & -1 \\ \vdots & \vdots \\ a_m^T & -1 \end{bmatrix}, \quad \tilde{b} = \begin{bmatrix} -b_1 \\ \vdots \\ -b_m \end{bmatrix}$$

# Sum of piecewise-linear functions

$$\text{minimize } f(x) + g(x) = \max_{i=1,\dots,m} (a_i^T x + b_i) + \max_{i=1,\dots,p} (c_i^T x + d_i)$$

**Cost function is piecewise-linear**

$$f(x) + g(x) = \max_{\substack{i=1,\dots,m \\ j=1,\dots,p}} ((a_i + c_j)^T x + (b_i + d_j))$$

**Equivalent linear optimization**

$$\text{minimize } t_1 + t_2$$

$$\text{subject to } a_i^T x + b_i \leq t_1, \quad i = 1, \dots, m$$

$$c_i^T x + d_i \leq t_2, \quad i = 1, \dots, p$$

**Matrix**

**notation?**

# Examples

# Cheapest cat food problem

- Choose quantities  $x_1, \dots, x_n$  of  $n$  ingredients each with unit cost  $c_j$ .
- Each ingredient  $j$  has nutritional content  $a_{ij}$  for nutrient  $i$ .
- Require a minimum level  $b_i$  for each nutrient  $i$ .

$$\begin{aligned} & \text{minimize} && \sum_{j=1}^n c_j x_j \\ & \text{subject to} && \sum_{j=1}^n a_{ij} x_j \geq b_i, \quad i = 1 \dots m \\ & && x_j \geq 0, \quad j = 1 \dots n \end{aligned}$$



[Photo Phoebe, my cat]


**Would you give her  
the optimal food ?**

# Data-fitting example

Fit a linear function  $f(z) = a + cz$  to  $m$  data points  $(z_i, f_i)$ :

Approximation problem  $Ax \approx b$  where


$$\underbrace{\begin{bmatrix} 1 & z_1 \\ \vdots & \vdots \\ 1 & z_m \end{bmatrix}}_A \underbrace{\begin{bmatrix} a \\ c \end{bmatrix}}_x \approx \underbrace{\begin{bmatrix} f_1 \\ \vdots \\ f_m \end{bmatrix}}_b$$



# Data-fitting example

Fit a linear function  $f(z) = a + cz$  to  $m$  data points  $(z_i, f_i)$ :

Approximation problem  $Ax \approx b$  where



$$\underbrace{\begin{bmatrix} 1 & z_1 \\ \vdots & \vdots \\ 1 & z_m \end{bmatrix}}_A \underbrace{\begin{bmatrix} a \\ c \end{bmatrix}}_x \approx \underbrace{\begin{bmatrix} f_1 \\ \vdots \\ f_m \end{bmatrix}}_b$$

**Least squares way:**


$$\text{minimize } \sum_{i=1}^m (Ax - b)_i^2 = \|Ax - b\|_2^2$$

**Good news:** solution is in closed form  $x^* = (A^T A)^{-1} A^T b$   
**Bad news:** solution is very sensitive to outliers!

# Data-fitting example

Fit a linear function  $f(z) = a + cz$  to  $m$  data points  $(z_i, f_i)$ :

Approximation problem  $Ax \approx b$  where



$$\underbrace{\begin{bmatrix} 1 & z_1 \\ \vdots & \vdots \\ 1 & z_m \end{bmatrix}}_A \underbrace{\begin{bmatrix} a \\ c \end{bmatrix}}_x \approx \underbrace{\begin{bmatrix} f_1 \\ \vdots \\ f_m \end{bmatrix}}_b$$

**A different way:**

$$\text{minimize } \sum_{i=1}^m |Ax - b|_i = \|Ax - b\|_1$$

**Good news:** solution is much more robust to outliers.

**Bad news:** there is no closed form solution.

# 1-norm approximation

$$\text{minimize} \quad \|Ax - b\|_1$$

The **1-norm** of  $m$ -vector  $y$  is

$$\|y\|_1 = \sum_{i=1}^m |y_i| = \sum_{i=1}^m \max\{y_i, -y_i\}$$

## Equivalent problem

$$\text{minimize} \quad \sum_{i=1}^m u_i$$

$$\text{subject to} \quad -u \leq Ax - b \leq u$$

## Matrix notation



$$\begin{bmatrix} 0 \\ 1 \end{bmatrix}^T \begin{bmatrix} x \\ u \end{bmatrix}$$

$$\begin{bmatrix} A & -I \\ -A & -I \end{bmatrix} \begin{bmatrix} x \\ u \end{bmatrix} \leq \begin{bmatrix} b \\ -b \end{bmatrix}$$

# Comparison with least-squares

Histogram of residuals  $Ax - b$  with randomly generated  $A \in \mathbf{R}^{200 \times 80}$

$$x_{\text{ls}} = \operatorname{argmin} \|Ax - b\|, \quad x_{\ell_1} = \operatorname{argmin} \|Ax - b\|_1$$



$\ell_1$ -norm distribution is **wider** with a **high peak at zero**

# $\ell_\infty$ -norm (Chebyshev) approximation

$$\text{minimize} \quad \|Ax - b\|_\infty$$

The  $\infty$ -norm of  $m$ -vector  $y$  is

$$\|y\|_\infty = \max_{i=1,\dots,m} |y_i| = \max_{i=1,\dots,m} \max\{y_i, -y_i\}$$

## Equivalent problem

$$\begin{aligned} & \text{minimize} && t \\ & \text{subject to} && -t\mathbf{1} \leq Ax - b \leq t\mathbf{1} \end{aligned}$$

## Matrix notation

$$\begin{aligned} & \text{minimize} && \begin{bmatrix} 0 \\ 1 \end{bmatrix}^T \begin{bmatrix} x \\ t \end{bmatrix} \\ & \text{subject to} && \begin{bmatrix} A & -1 \\ -A & -1 \end{bmatrix} \begin{bmatrix} x \\ t \end{bmatrix} \leq \begin{bmatrix} b \\ -b \end{bmatrix} \end{aligned}$$

# Sparse signal recovery via $\ell_1$ -norm minimization

$\hat{x} \in \mathbf{R}^n$  is unknown signal, known to be sparse

We make linear measurements  $y = A\hat{x}$  with  $A \in \mathbf{R}^{m \times n}$ ,  $m < n$

Estimate signal with smallest  $\ell_1$ -norm, consistent with measurements

$$\begin{aligned} &\text{minimize} && \|x\|_1 \\ &\text{subject to} && Ax = y \end{aligned}$$

## Equivalent linear optimization

$$\begin{aligned} &\text{minimize} && \mathbf{1}^T u \\ &\text{subject to} && -u \leq x \leq u \\ & && Ax = y \end{aligned}$$


# Sparse signal recovery via $\ell_1$ -norm minimization

## Example

Exact signal  $\hat{x} \in \mathbf{R}^{1000}$

10 nonzero components


Random  $A \in \mathbf{R}^{100 \times 1000}$



The least squares estimate  
cannot recover the sparse signal

minimize  $\|x\|_2^2$


subject to  $Ax = y$



The  $\ell_1$ -norm estimate is **exact**

minimize  $\|x\|_1$

subject to  $Ax = y$



# Sparse signal recovery via $\ell_1$ -norm minimization

## Exact recovery

When are these two problems equivalent?

$$\text{minimize} \quad \text{card}(x)$$

$$\text{subject to} \quad Ax = y$$

$$\text{minimize} \quad \|x\|_1$$

$$\text{subject to} \quad Ax = y$$

$\text{card}(x)$  is cardinality (number of nonzero components) of  $x$

We say  $A$  allows **exact recovery** of  $k$ -sparse vectors if

$$\hat{x} = \underset{Ax=y}{\operatorname{argmin}} \|x\|_1 \quad \text{when } y = A\hat{x} \text{ and } \text{card}(\hat{x}) \leq k$$

It depends on the nullspace<sup>1</sup> of the “measurement matrix”  $A$

# Linear classification


## Support vector machine (linear separation)

Given a set of points  $\{v_1, \dots, v_N\}$  with binary labels  $s_i \in \{-1, 1\}$

Find hyperplane that strictly separates the two classes

$$a^T v_i + b > 0 \quad \text{if} \quad s_i = 1$$

$$a^T v_i + b < 0 \quad \text{if} \quad s_i = -1$$



Homogeneous in  $(a, b)$ , hence equivalent to the linear inequalities (in  $a, b$ )

$$s_i(a^T v_i + b) \geq 1$$

# Linear classification

## Separable case

### Feasibility problem

find

$a, b$

subject to

$s_i(a^T v_i + b) \geq 1, \quad i = 1, \dots, N$

Which can be seen as a special case of LP with

minimize

0

subject to

$s_i(a^T v_i + b) \geq 1, \quad i = 1, \dots, N$

$p^* = 0$  if problem feasible (points separable)

$p^* = \infty$  if problem infeasible (points not separable) —————> **What then?**


# Linear classification

## Approximate linear separation of non-separable points

$$\text{minimize} \quad \sum_{i=1}^N (1 - s_i(a^T v_i + b))_+ = \sum_{i=1}^N \max\{0, 1 - s_i(a^T v_i + b)\}$$

If  $v_i$  misclassified,  $1 - s_i(a^T v_i + b)$  is the penalty

Piecewise-linear minimization problem with variables  $a, b$



# Linear classification

## Approximate linear separation of non-separable points


$$\text{minimize} \quad \sum_{i=1}^N \max\{0, 1 - s_i(a^T v_i + b)\}$$

### Equivalent problem

$$\text{minimize} \quad \sum_{i=1}^N u_i$$

$$\text{subject to} \quad 1 - s_i(v_i^T a + b) \leq u_i, \quad i = 1, \dots, N$$

$$0 \leq u_i, \quad i = 1, \dots, N$$



### Matrix notation?

# Modelling software for linear programs

Modelling tools simplify the formulation of LPs (and other problems)

- Accept optimization problem in common notation ( $\max$ ,  $\|\cdot\|_1, \dots$ )
- Recognize problems that can be converted to LPs
- Express the problem in input format required by a specific LP solver

## Examples

- AMPL, GAMS
- CVX, YALMIP (Matlab)
- CVXPY, Pyomo (Python)
- JuMP.jl, Convex.jl (Julia)

# CVXPY example

$$\begin{aligned} & \text{minimize} && \|Ax - b\|_1 \\ & \text{subject to} && 0 \leq x \leq 1 \end{aligned}$$

```
x = cp.Variable(n)
objective = cp.Minimize(cp.norm(A*x - b, 1))
constraints = [0 <= x, x <= 1]
problem = cp.Problem(objective, constraints)

# The optimal objective value is returned by `problem.solve()`.

result = problem.solve()

# The optimal value for x is stored in `x.value`.
print(x.value)
```

# Why linear optimization?

## “Easy” to solve

- It is solvable in polynomial time, and it is tractable in practice
- State-of-the-art software can solve LPs with tens of thousands of variables.  
We can solve LPs with millions of variables with specific structure.

## Extremely versatile

It can model many real-world problems, either exactly or approximately.

## Fundamental

The theory of linear optimization lays the foundation for most optimization theories

# Next lecture

## Geometry of linear optimization

- Polyhedra
- Extreme points
- Basic feasible solutions